Slide 1

Uncertain Knowledge and Reasoning

Outline

- Acting under uncertainty
- Decision-theoretic agent
- Probability
- Bayes' rule

Slide 2

Acting under Uncertainty

- Agents almost never have access to the whole truth about their environment
- Qualification problem:
  how to define the circumstances under which a given action is *guaranteed* to work.
- Rational decision
  depends on the relative importance of the various goals and the likelihood that, and degree to which, they will be achieved.

Slide 3

Handling uncertain knowledge

- Example: diagnostic knowledge
  1. $\forall p(Symptom(p, Toothache) \rightarrow Disease(p, Cavity))$
  2. $\forall p(Symptom(p, Toothache) \rightarrow$
     $Disease(p, Cavity) \vee Disease(p, GumDisease) \vee$
     $Disease(p, ImpactedWisdom) \vee \cdots)$
  3. $\forall p(Disease(p, Cavity)) \rightarrow Symptom(p, Toothache)$
- Difficulties
  —Laziness
  —Theoretical ignorance
  —Practical ignorance

Slide 4

Probability theory

- Agent's knowledge: degree of belief
- Probability: assigns a degree of belief between 0 and 1 to sentences.
  A way of summarizing the uncertainty.
- Sentences either true or false.
  — A probability of 0.8 that cavity if toothache.
  — A probability of 0 $\mapsto$ false
  — A probability of 1 $\mapsto$ true
- Degree of belief vs. degree of truth
  Probability theory vs. fuzzy logic.

Slide 5

- The probability that an agent assigns to a proposition depends of the percepts (evidence)
- Analogous to logical consequence
- Prior/unconditional probability
- Posterior/conditional probability

Slide 6

Decision-theoretic agent

- Utility theory (preferences)
- Decision theory = probability theory + utility theory
  Maximum expected utility: An agent is rational iff it chooses the action that yields the highest expected utility, averaged over all the possible outcomes of the action.

```
function DT-AGENT(percept) returns an action
  static: a set probabilistic beliefs about the state of the world

  calculate updated probabilities for current state based on
      available evidence including current percept and previous action
  calculate outcome probabilities for actions,
      given action descriptions and probabilities of current states
  select action with highest expected utility
      given probabilities of outcomes and utility information
  return action
```


Slide 7

Probability

- Prior/unconditional probability: $P(Cavity) = 0.1$
- Random variables
  $P(Weather = Sunny) = 0.7 \quad P(Cavity = true) = 0.1$
  $P(Weather = Rain) = 0.2 \quad P(Cavity = false) = 0.9$
  $P(Weather = Cloudy) = 0.08$
  $P(Weather = Snow) = 0.02$
- Probability distribution:
  $\mathbf{P}(Weather) = \langle 0.7, 0.2, 0.08, 0.02 \rangle$
  $\mathbf{P}(Weather, Cavity)$: 4×2 table

Slide 8

Conditional/posterior probability

$P(Cavity|Toothache) = 0.8$

$$P(A|B) = \frac{P(A \wedge B)}{P(B)}$$

whenever $P(B) > 0$

- Product rule: $P(A \wedge B) = P(A|B)P(B)$
  $P(A \wedge B) = P(B|A)P(A)$
- $\mathbf{P}(X \wedge Y) = \mathbf{P}(X|Y)\mathbf{P}(Y)$
  $\Rightarrow$
  $P(X = x_1 \wedge Y = y_2) = P(X = x_1|Y = y_2)P(Y = y_2))$

Slide 9

Axioms of probability

1. $0 \leq P(A) \leq 1$
2. $P(\phi) = 0/1$ if ϕ is unsatisfiable/valid
3. $P(A \vee B) = P(A) + P(B) - P(A \wedge B)$



Slide 10

- $P(A \wedge B) = P(A) + P(B) - P(A \vee B)$ (by 3)
- $P(\neg A) = 1 - P(A)$

$$\begin{aligned} P(A \vee \neg A) &= P(A) + P(\neg A) - P(A \wedge \neg A) \text{ (by 3)} \\ 1 &= P(A) + P(\neg A) - 0 \text{ (by 2)} \\ P(\neg A) &= 1 - P(A) \end{aligned}$$

Note that $A \vee \neg A$ is valid and $A \wedge \neg A$ is unsatisfiable.

Slide 11

Why the axioms of probability are reasonable

(Bruno de Finetti, 1931):
If Agent 1 expresses a set of degrees of belief that violate the axioms of probability theory then there is a betting strategy for Agent 2 that guarantees that Agent 1 will lose money.

Example.

Agent 1		Agent 2		Outcome for Agent 1			
Proposition	Belief	Bet	Stakes	$A \wedge B$	$A \wedge \neg B$	$\neg A \wedge B$	$\neg A \wedge \neg B$
A	0.4	A	4 to 6	-6	-6	4	4
B	0.3	B	3 to 7	-7	3	-7	3
$A \vee B$	0.8	$\neg(A \vee B)$	2 to 8	2	2	2	-8
				-11	-1	-1	-1

Slide 12

The joint probability distribution

- Atomic event: assignment of particular values to all the variables.
- The joint probability distribution $\mathbf{P}(X_1, \ldots, X_n)$: assigns probabilities to all possible atomic events.

	$Toothache$	$\neg Toothache$
$Cavity$	0.04	0.06
$\neg Cavity$	0.01	0.89

Slide 13

- Entries in the table sum to 1
- Can be used to compute any probabilistic statement

$P(Cavity) = 0.04 + 0.06 = 0.10$

$P(Cavity \vee Toothache) = 0.04 + 0.06 + 0.01 = 0.11$

$$\begin{aligned} P(Cavity|Toothache) &= \frac{P(Cavity \wedge Toothache)}{P(Toothache)} \\ &= \frac{0.04}{0.04 + 0.01} = 0.80 \end{aligned}$$

- For n Boolean variables, 2^n entries.

Slide 14

Bayes' Rule

- Bayes' rule is obtained from the product rule
  ($P(A \wedge B) = P(A|B)P(B)$, $P(A \wedge B) = P(B|A)P(A)$)

$$P(B|A) = \frac{P(A|B)P(B)}{P(A)}$$

- For multi-valued variables:

$$\mathbf{P}(Y|X) = \frac{\mathbf{P}(X|Y)\mathbf{P}(Y)}{\mathbf{P}(X)}$$

- For background evidence E:

$$\mathbf{P}(Y|X, E) = \frac{\mathbf{P}(X|Y, E)\mathbf{P}(Y|E)}{\mathbf{P}(X|E)}$$

Slide 15

Applying Bayes's rule

$$\begin{aligned} P(S|M) &= 0.5 \\ P(M) &= 1/50000 \\ P(S) &= 1/20 \\ P(M|S) &= \frac{P(S|M)P(M)}{P(S)} \\ &= \frac{0.5 \times 1/50000}{1/20} = 0.0002 \end{aligned}$$

Diagnostic knowledge is often more tenuous than causal knowledge.

Slide 16

Normalization

How to avoid direct assessment of the prior probabilities of the "symptoms"?

$$\begin{aligned} P(M|S) &= \frac{P(S|M)P(M)}{P(S)} \\ P(\neg M|S) &= \frac{P(S|\neg M)P(\neg M)}{P(S)} \\ 1 &= P(M|S) + P(\neg M|S) \\ P(S) &= P(S|M)P(M) + P(S|\neg M)P(\neg M) \\ P(M|S) &= \frac{P(S|M)P(M)}{P(S|M)P(M) + P(S|\neg M)P(\neg M)} \end{aligned}$$

Slide 17

Combining evidence

$P(Cavity|Toothache) = 0.8$
$P(Cavity|Catch) = 0.95$

$$P(Cavity|Catch \wedge Toothache) = \frac{P(Catch \wedge Toothache|Cavity)P(Cavity)}{P(Catch \wedge Toothache)}$$

Many numbers have to be specified!
☞ Bayesian updating

$$P(Cavity|Toothache) = P(Cavity)\frac{P(Toothache|Cavity)}{P(Toothache)}$$

Slide 18

$P(Cavity|Toothache \wedge Catch) =$

$$P(Cavity|Toothache)\frac{P(Catch|Toothache \wedge Cavity)}{P(Catch|Toothache)} =$$
$$P(Cavity)\frac{P(Toothache|Cavity)}{P(Toothache)}\frac{P(Catch|Toothache \wedge Cavity)}{P(Catch|Toothache)}$$

Many numbers still!
☞ Conditional independence of $Tootache$ and $Catch$ given $Cavity$:
$P(Catch|Toothache \wedge Cavity) = P(Catch|Cavity)$
$P(Toothache|Catch \wedge Cavity) = P(Toothache|Cavity)$

Slide 19

- $P(Cavity|Toothache \wedge Catch) = P(Cavity)\frac{P(Toothache|Cavity)}{P(Toothache)}\frac{P(Catch|Cavity)}{P(Catch|Toothache)}$
- Provided we assess $P(Toothache|\neg Cavity)$ and $P(Catch|\neg Cavity)$
  $P(Toothache)P(Catch|Toothache)$ can be eliminated by normalization.
- Conditional independence information crucial.

Slide 20

Where do the probabilities come from?

- Frequentist
- Objectivist
- Subjectivist
- Reference class problem