


**Solutions to demonstration problems****Solution to Problem 4**

We transform the propositions into CNF and clauses. The last proposition in the table is the negation of statement “both red lights are not on at the same time”, that is,

$$\neg(\neg(P1 \wedge P2)) \equiv P1 \wedge P2.$$

$Pi \vee Ki \vee Vi$	$\{Pi, Ki, Vi\}$
$Pi \rightarrow \neg Ki \wedge \neg Vi \equiv \neg Pi \vee (\neg Ki \wedge \neg Vi)$ $\equiv (\neg Pi \vee \neg Ki) \wedge (\neg Pi \vee \neg Vi)$	$\{\neg Pi, \neg Ki\}, \{\neg Pi, \neg Vi\}$
$Ki \rightarrow \neg Pi \wedge \neg Vi \equiv (\neg Ki \vee \neg Pi) \wedge (\neg Ki \vee \neg Vi)$	$\{\neg Pi, \neg Ki\}, \{\neg Ki, \neg Vi\}$
$Vi \rightarrow \neg Pi \wedge \neg Ki \equiv (\neg Vi \vee \neg Pi) \wedge (\neg Vi \vee \neg Ki)$	$\{\neg Pi, \neg Vi\}, \{\neg Ki, \neg Vi\}$
$\neg(V1 \wedge V2) \equiv \neg V1 \vee \neg V2$	$\{\neg V1, \neg V2\}$
$P1 \rightarrow (K2 \vee V2) \equiv \neg P1 \vee K2 \vee V2$	$\{\neg P1, K2, V2\}$
$P2 \rightarrow (K1 \vee V1) \equiv \neg P2 \vee K1 \vee V1$	$\{\neg P2, K1, V1\}$
$P1 \wedge P2$	$\{P1\}, \{P2\}$

We show that the set of clauses given in the table is unsatisfiable (empty clause  $\square$  means contradiction), which implies that  $\neg(P1 \wedge P2)$  is derivable from the other clauses.



### Solution to Problem 5

The chemical reactions can be formalized as implications, which can then be transformed into clausal form. The resulting clauses are:

(1)

$$\begin{aligned} & \text{MgO} + \text{H}_2 \rightarrow \text{Mg} + \text{H}_2\text{O} \\ \implies & \text{MgO} \wedge \text{H}_2 \rightarrow \text{Mg} \wedge \text{H}_2\text{O} \\ \implies & \neg \text{MgO} \vee \neg \text{H}_2 \vee (\text{Mg} \wedge \text{H}_2\text{O}) \\ \implies & (\neg \text{MgO} \vee \neg \text{H}_2 \vee \text{Mg}) \wedge (\neg \text{MgO} \vee \neg \text{H}_2 \vee \text{H}_2\text{O}) \end{aligned}$$

The first reaction results in two clauses:  $\{\neg \text{MgO}, \neg \text{H}_2, \text{Mg}\}$  and  $\{\neg \text{MgO}, \neg \text{H}_2, \text{H}_2\text{O}\}$ .

(2)

$$\begin{aligned} & \text{C} + \text{O}_2 \rightarrow \text{CO}_2 \\ \implies & \text{C} \wedge \text{O}_2 \rightarrow \text{CO}_2 \\ \implies & \neg \text{C} \vee \neg \text{O}_2 \vee \text{CO}_2 \\ \implies & \{\neg \text{C}, \neg \text{O}_2, \text{CO}_2\} \end{aligned}$$


(3)

$$\begin{aligned} & \text{CO}_2 + \text{H}_2\text{O} \rightarrow \text{H}_2\text{CO}_3 \\ \implies & \text{CO}_2 \wedge \text{H}_2\text{O} \rightarrow \text{H}_2\text{CO}_3 \\ \implies & \neg \text{CO}_2 \vee \neg \text{H}_2\text{O} \vee \text{H}_2\text{CO}_3 \\ \implies & \{\neg \text{CO}_2, \neg \text{H}_2\text{O}, \text{H}_2\text{CO}_3\} \end{aligned}$$

The elements available at the start are:

$$\begin{aligned} & \text{MgO} \wedge \text{H}_2 \wedge \text{O}_2 \wedge \text{C} \\ \implies & \{\text{MgO}\}, \{\text{H}_2\}, \{\text{O}_2\}, \{\text{C}\} \end{aligned}$$

We denote the above set of clauses with  $\Sigma$ . now we want to prove that  $\Sigma \models \text{H}_2\text{CO}_3$ . The proof is constructed by showing that  $\Sigma \cup \{\neg \text{H}_2\text{CO}_3\}$  is unsatisfiable.



### Solution to Problem 6

The solution is obtained from “Computational Complexity” by C. Papadimitriou. A deterministic Turing machine is a quadruple  $\langle A, S, s_0, t \rangle$ , where

- $A$  is the alphabet,
- $S$  is the set of states,
- $t : S \times A \rightarrow S \times A \times \{\rightarrow, \leftarrow, \downarrow\}$  is the state transition function
- $s_0 \in S$  is the start state.

For our machine we have  $S = \{s\}$ ,  $A = \{0, 1\}$ ,  $s_0 = s$  and the state transition function is given in the following table:

$p \in S$	$\sigma \in A$	$t(p, \sigma)$
$s$	0	$(h, 1, -)$
$s$	1	$(s, 0, \rightarrow)$
$s$	$\sqcup$	$(h, 1, -)$
$s$	$\triangleright$	$(s, \triangleright, \rightarrow)$

With input 1101 the computation goes as follows:  $(s, \triangleright, 1101) \xrightarrow{M} (s, \triangleright 0, 101) \xrightarrow{M} (s, \triangleright 00, 01) \xrightarrow{M} (h, \triangleright 001, 1)$ .

**Solution to Problem 7**

The problem of 3-coloring a graph is as follows: “give a graph  $G$ , is there a way to color the nodes in  $G$  using 3 colors so that no two adjacent nodes have same color?”

Let  $N = \{n_1, n_2, \dots, n_m\}$  be the set of nodes and  $E \subseteq N \times N$  the set of edges.

For each node  $n_i$  we take atomic propositions  $R_{n_i}, G_{n_i}, B_{n_i}$  to denote that node  $n_i$  is colored red, green or blue, respectively.

Each node is colored with some color, that is,  $R_{n_i} \vee G_{n_i} \vee B_{n_i}$ , for each  $n_i$ .

No node is colored with two different colors, that is,

$$(R_{n_i} \rightarrow (\neg G_{n_i} \wedge \neg B_{n_i})) \wedge (G_{n_i} \rightarrow (\neg R_{n_i} \wedge \neg B_{n_i})) \wedge (B_{n_i} \rightarrow (\neg R_{n_i} \wedge \neg G_{n_i})),$$

for each  $n_i$ .

Finally, two adjacent color can't have same color, that is,

$$(R_n \rightarrow \neg R_m) \wedge (G_n \rightarrow \neg G_m) \wedge (B_n \rightarrow \neg B_m),$$

for each  $(n, m) \in E$ .

Now, if we take the conjunction of all these propositions (denoted by  $\phi$ ), then  $\phi$  is satisfiable iff the graph has a 3-coloring (the proof is omitted).