

T-79.5102

Syksy 2005

Laskennallisen logiikan erikoiskurssi

Laskuharjoitus 7

Ratkaisut

1. a)

$$\begin{aligned}T_S(\emptyset) \uparrow 0 &= \{R\} \\T_S(\emptyset) \uparrow 1 &= \{R, Q_1\} \\T_S(\emptyset) \uparrow 2 &= \{R, Q_1, P_1\} \\T_S(\emptyset) \uparrow 3 &= \{R, Q_1, P_1, S_1\} \\T_S(\emptyset) \uparrow 4 &= \{R, Q_1, P_1, S_1\}\end{aligned}$$

Siis $M_S = \{R, Q_1, P_1, S_1\}$.

b)

$$\begin{array}{ll}R & [R] \\Q_1 & [Q_1 \leftarrow R] \\P_1 & [P_1 \leftarrow Q_1] \\S_1 & [S_1 \leftarrow P_1 \wedge Q_1]\end{array}$$

c)

$$\begin{aligned}T_S(\emptyset) \uparrow 0 &= \{Q(a)\} \\T_S(\emptyset) \uparrow 1 &= \{Q(g^i(a))\}_{i=0}^1 \\T_S(\emptyset) \uparrow 2 &= \{Q(g^i(a))\}_{i=0}^2 \\&\vdots \\T_S(\emptyset) \uparrow n &= \{Q(g^i(a))\}_{i=0}^n \\&\vdots \\T_S(\emptyset) \uparrow \omega &= \{Q(g^i(a))\}_{i=0}^\infty\end{aligned}$$

2. a) Klausuulijoukon S Herbrand-instantiaatio S_H on:

$$\begin{aligned} & \{ d(1) \leftarrow a(1), b(1); d(2) \leftarrow a(2), b(2); \\ & \quad d(3) \leftarrow a(3), b(3); d(4) \leftarrow a(4), b(4); \\ & \quad e(1, 1) \leftarrow d(1), d(1); e(1, 2) \leftarrow d(1), d(2); \\ & \quad e(1, 3) \leftarrow d(1), d(3); e(1, 4) \leftarrow d(1), d(4); \\ & \quad e(2, 1) \leftarrow d(2), d(1); e(2, 2) \leftarrow d(2), d(2); \\ & \quad e(2, 3) \leftarrow d(2), d(3); e(2, 4) \leftarrow d(2), d(4); \\ & \quad e(3, 1) \leftarrow d(3), d(1); e(3, 2) \leftarrow d(3), d(2); \\ & \quad e(3, 3) \leftarrow d(3), d(3); e(3, 4) \leftarrow d(3), d(4); \\ & \quad e(4, 1) \leftarrow d(4), d(1); e(4, 2) \leftarrow d(4), d(2); \\ & \quad e(4, 3) \leftarrow d(4), d(3); e(4, 4) \leftarrow d(4), d(4); \\ & \quad a(1); a(2); b(2); b(3); c(4) \} \end{aligned}$$

b) Ja sen minimimalli:

$$\begin{aligned} T_{S_H}(\emptyset) \uparrow 0 &= \{a(1), a(2), b(2), b(3), c(4)\} \\ T_{S_H}(\emptyset) \uparrow 1 &= \{d(2), a(1), a(2), b(2), b(3), c(4)\} \\ T_{S_H}(\emptyset) \uparrow 3 &= \{e(2, 2), d(2), a(1), a(2), b(2), b(3), c(4)\} \end{aligned}$$

c) Pienin instantiaatio G saadaan S_H :sta jättämällä pois kaikki säännöt, joiden vartalossa esiintyy totutumattomia atomeita:

$$G = \{ d(2) \leftarrow a(2), b(2), e(2, 2) \leftarrow d(2), d(2); \\ a(1); a(2); b(2); b(3); c(4) \}$$