

## Combinatorial Models and Stochastic Algorithms

## Tutorial 3, February 8

## Problems

1. Denote  $C = \{1, \dots, c\}$ , and let  $\pi$  be any probability distribution on the state set  $S = C^n$ . Prove that the basic Gibbs sampler for  $\pi$  has  $\pi$  as its stationary distribution. (*Hint:* Generalise the argument used in the lecture notes in the case of the Gibbs sampler for the hard-core model.)
2. Consider an arbitrary distribution  $\pi$  on the state set  $S = \{0, 1\}^n$ . Design for  $\pi$  both (a) a basic Gibbs sampler, and (b) a Metropolis sampler using the Hamming neighbourhood, where  $S$  is viewed as a graph whose two nodes are neighbours if and only if their co-ordinate vectors differ in exactly one position. Are the two samplers the same?
3. Verify the claims in Proposition 3.5 of the lecture notes. That is: given a regular reversible Markov chain  $\mathcal{M}$  on state set  $S = \{1, \dots, n\}$  with transition matrix  $P$  and stationary distribution  $\pi$ , show that the chain  $\mathcal{M}'$  with transition matrix  $P' = \frac{1}{2}(I_n + P)$  is also regular and reversible, has same stationary distribution  $\pi$  as  $\mathcal{M}$ , its eigenvalues satisfy  $1 = \lambda'_1 > \lambda'_2 \geq \dots \geq \lambda'_n > 0$ , and  $\lambda'_{\max} = \lambda'_2 = \frac{1}{2}(1 + \lambda_2)$ , where  $\lambda_2$  is the second largest eigenvalue of  $\mathcal{M}$ . Estimate the effect of the change from  $P$  to  $P'$  on the convergence rate of the chain.
4. Consider a random walk on an undirected graph  $G = (V, E)$ , where transitions are made from each node  $u$  to an adjacent node with uniform probability  $\beta/d$ , where  $d$  is the maximum degree of any node in  $G$  and  $\beta \leq 1$  is a positive constant. In addition, each node  $u$  has a self-loop probability of  $1 - \beta \deg(u)/d$ . Prove that if  $G$  is connected and  $\beta < 1$ , then the corresponding Markov chain  $\mathcal{M}_G$  is regular and reversible, with uniform stationary distribution. Moreover, show that the conductance of  $\mathcal{M}_G$  is given by the formula

$$\Phi = \beta\mu(G)/d,$$

where  $\mu(G)$  is the *edge magnification* (also known as the *isoperimetric number* or *Cheeger constant*) of  $G$ , defined as

$$\mu(G) = \min_{0 < |U| \leq |V|/2} \frac{|\partial(U)|}{|U|},$$

where  $\partial(U) = \{\{u, v\} \in E \mid u \in U, v \notin U\}$ .

5. Based on the result of Problem 4, calculate an upper bound on the mixing time of a simple symmetric random walk on an  $n \times n$  square lattice with self-loop parameter  $0 < 1 - \beta < 1$  and periodic boundary conditions (i.e. each node  $(i, j)$ ,  $i, j = 0, \dots, n - 1$ , has as neighbours the nodes  $(i \pm 1, j \pm 1) \bmod n$ ).